

BROILER

YIELD PLUS x ROSS 308 AP

Performance Objectives

Revised 2021



Introduction

This booklet contains the performance objectives for the Yield Plus® (YP) x Ross® 308 AP and is to be used with the **Ross Broiler Management Handbook**.

Performance

These objectives indicate the performance achievable under good management and environmental conditions and when feeding recommended nutrient levels.

Producers may find that local factors prevent such performance being achieved. For example:

- The availability of raw materials may limit nutrient content and intake.
- Extreme climatic conditions will reduce performance.
- Economic considerations may limit choice of production systems.

Therefore, average performance may be lower than the figures presented here.

The objectives are presented in two sections to reflect the global nature of the publication.

Section 1 g contains the performance data in metric measurements and

Section 2 lb contains imperial measurements.

In the tables values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.

Yields will vary between processing plants depending on type of equipment used (e.g. carcass chilling technology, automated versus manual de-boning) and the exact portion being produced.

Every attempt has been made to ensure the accuracy and relevance of the information presented, however, Aviagen® accepts no liability for the consequences of using this information for the management of chickens.

For further information on the management of Ross stock, please contact your local Ross representative.

Contents

02		Key Management Points
03	Section 1 <i>g</i>	As-Hatched Performance
04	Section 1 <i>g</i>	Male Performance
05	Section 1 <i>g</i>	Female Performance
07	Section 2 <i>lb</i>	As-Hatched Performance
08	Section 2 <i>lb</i>	Male Performance
09	Section 2 <i>lb</i>	Female Performance
11		Carcass Yield

Key Management Points

Cost effective production of chicken meat depends on achieving good bird performance and the following points are important for optimizing performance of the YP x Ross 308 AP broiler:

- Maximize chick quality by good management of hatching, storage and transport conditions.
- Design the brooding set-up to ensure easy access to water and feed at placement, and to ease the transition between supplementary systems and the automated feeders and drinkers at 4-5 days. Feed a highly digestible and nutritionally balanced Starter diet.
- Keep chicks in their thermal comfort zone by monitoring chick behavior, but beware of low relative humidities (less than 50% RH). Establish a minimum ventilation program from day one.
- Monitor crop fill, feeding and drinking behavior and 7-day live weight to allow continuous improvement of the brooding set-up.
- Keep birds in their thermal comfort zone throughout the growing period. Fast growing broilers produce large amounts of heat, particularly in the second half of the grow-out period. Keeping ambient temperatures less than 21°C (69.8°F) from 21 days onwards may improve growth rates.
- Maintain high standards of biosecurity and cleanliness to keep disease to a minimum.

As-Hatched Performance

Day	Body weight (g) ¹	Daily gain (g)	Av. daily gain/week (g)	Daily intake (g)	Cum. intake (g) ²	FCR ³
0	44					
1	60	16			11	0.183
2	77	17		15	26	0.336
3	96	19		19	44	0.463
4	118	22		23	67	0.570
5	142	25		27	94	0.659
6	170	28		31	125	0.735
7	201	31	22.42	35	160	0.799
8	234	34		40	200	0.854
9	271	37		44	244	0.900
10	312	40		49	293	0.941
11	356	44		54	347	0.976
12	403	47		59	406	1.007
13	454	51		64	470	1.035
14	508	54	43.94	69	538	1.060
15	566	58		74	612	1.083
16	627	61		79	692	1.104
17	691	64		85	776	1.124
18	758	67		90	866	1.143
19	829	71		95	961	1.161
20	902	74		101	1062	1.178
21	979	76	67.23	106	1169	1.195
22	1058	79		112	1280	1.211
23	1139	82		117	1398	1.228
24	1223	84		123	1521	1.244
25	1310	86		129	1650	1.260
26	1398	88		134	1784	1.277
27	1489	90		140	1923	1.293
28	1581	92	86.00	145	2068	1.309
29	1674	94		150	2219	1.326
30	1769	95		156	2375	1.343
31	1866	96		161	2536	1.360
32	1963	97		166	2702	1.377
33	2062	98		171	2873	1.395
34	2161	99		176	3050	1.413
35	2260	100	97.09	181	3231	1.431
36	2360	100		186	3417	1.449
37	2461	100		191	3607	1.467
38	2561	101		195	3802	1.486
39	2662	101		199	4002	1.505
40	2762	100		204	4205	1.524
41	2862	100		208	4413	1.544
42	2962	100	100.25	212	4625	1.563
43	3061	99		215	4840	1.583
44	3160	99		219	5059	1.603
45	3258	98		223	5282	1.623
46	3355	97		226	5508	1.644
47	3451	96		229	5737	1.665
48	3547	95		232	5969	1.686
49	3641	94	97.00	235	6204	1.707
50	3734	93		238	6442	1.728
51	3826	92		240	6683	1.749
52	3917	91		243	6926	1.771
53	4007	90		245	7171	1.793
54	4095	88		247	7418	1.815
55	4182	87		249	7668	1.837
56	4267	86	89.48	251	7919	1.859
57	4351	84		253	8172	1.882
58	4434	83		255	8427	1.904
59	4515	81		256	8683	1.927
60	4595	80		258	8941	1.950
61	4673	78		259	9200	1.973
62	4750	77		260	9460	1.996
63	4825	75	79.67	261	9721	2.019
64	4899	74		262	9983	2.042
65	4971	72		263	10246	2.066
66	5041	71		264	10510	2.089
67	5111	69		264	10774	2.113
68	5178	68		265	11039	2.137
69	5244	66		265	11304	2.161
70	5309	65	69.09	266	11570	2.185

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table the values are rounded. This may result in small inaccuracies when using the objectives to calculate other performance statistics.

Male Performance

Day	Body weight (g) ¹	Daily gain (g)	Av. daily gain/week (g)	Daily intake (g)	Cum. intake (g) ²	FCR ³
0	44					
1	60	16			10	0.169
2	77	17		14	24	0.312
3	97	20		18	42	0.433
4	119	22		22	64	0.536
5	144	25		26	90	0.624
6	172	28		30	120	0.699
7	204	31	22.85	35	155	0.764
8	238	35		40	195	0.819
9	276	38		45	240	0.868
10	318	42		50	289	0.910
11	363	45		55	344	0.947
12	412	49		60	404	0.979
13	465	53		65	469	1.009
14	521	56	45.41	71	540	1.035
15	581	60		76	616	1.059
16	645	64		82	698	1.082
17	712	67		88	785	1.102
18	783	71		93	879	1.122
19	857	74		99	978	1.140
20	935	77		105	1083	1.158
21	1015	81	70.58	111	1194	1.176
22	1099	84		117	1311	1.192
23	1186	87		123	1434	1.209
24	1275	89		129	1563	1.225
25	1367	92		135	1697	1.241
26	1462	94		141	1838	1.258
27	1559	97		147	1985	1.274
28	1657	99	91.70	153	2138	1.290
29	1758	101		159	2297	1.306
30	1861	103		165	2461	1.323
31	1965	104		170	2632	1.339
32	2070	105		176	2808	1.356
33	2177	107		182	2989	1.373
34	2284	108		187	3176	1.390
35	2393	109	105.10	192	3369	1.408
36	2502	109		198	3567	1.425
37	2612	110		203	3769	1.443
38	2722	110		208	3977	1.461
39	2833	110		213	4190	1.479
40	2943	110		218	4408	1.498
41	3053	110		222	4630	1.516
42	3163	110	110.06	227	4856	1.535
43	3273	110		231	5087	1.554
44	3383	109		235	5322	1.573
45	3491	109		239	5561	1.593
46	3600	108		243	5804	1.612
47	3707	107		247	6051	1.632
48	3813	106		250	6301	1.652
49	3919	105	107.90	253	6554	1.673
50	4023	104		257	6811	1.693
51	4127	103		260	7070	1.713
52	4229	102		263	7333	1.734
53	4330	101		265	7598	1.755
54	4429	100		268	7866	1.776
55	4527	98		270	8136	1.797
56	4624	97	100.75	272	8409	1.818
57	4719	95		275	8683	1.840
58	4813	94		277	8960	1.862
59	4906	92		278	9238	1.883
60	4996	91		280	9519	1.905
61	5085	89		282	9800	1.927
62	5173	88		283	10083	1.949
63	5259	86	90.72	284	10368	1.971
64	5343	84		286	10654	1.994
65	5426	83		287	10940	2.016
66	5507	81		288	11228	2.039
67	5587	79		289	11517	2.061
68	5665	78		289	11806	2.084
69	5741	76		290	12096	2.107
70	5815	75	79.49	290	12386	2.130

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table the values are rounded. This may result in small inaccuracies when using the objectives to calculate other performance statistics.

Female Performance

Day	Body weight (g) ¹	Daily gain (g)	Av. daily gain/week (g)	Daily intake (g)	Cum. intake (g) ²	FCR ³
0	44					
1	59	16			12	0.197
2	76	17		16	27	0.359
3	95	19		20	47	0.493
4	117	21		23	70	0.603
5	141	24		27	98	0.695
6	168	27		31	129	0.771
7	198	30	21.99	36	165	0.834
8	230	33		40	205	0.888
9	266	36		44	249	0.933
10	306	39		49	297	0.972
11	348	42		53	350	1.006
12	394	46		58	408	1.035
13	443	49		62	470	1.062
14	495	52	42.46	67	537	1.085
15	550	55		72	609	1.107
16	608	58		77	685	1.127
17	669	61		82	767	1.146
18	733	64		87	853	1.164
19	800	67		92	945	1.181
20	870	70		97	1042	1.197
21	942	72	63.88	102	1143	1.214
22	1016	74		107	1250	1.230
23	1093	77		112	1362	1.246
24	1172	79		117	1479	1.263
25	1252	81		122	1602	1.279
26	1335	82		127	1729	1.295
27	1419	84		132	1861	1.312
28	1504	85	80.30	137	1999	1.329
29	1591	87		142	2141	1.346
30	1678	88		147	2288	1.363
31	1767	89		152	2440	1.381
32	1856	89		156	2596	1.399
33	1946	90		161	2757	1.417
34	2037	90		166	2923	1.435
35	2128	91	89.09	170	3093	1.454
36	2219	91		174	3267	1.472
37	2310	91		178	3445	1.492
38	2400	91		182	3627	1.511
39	2491	91		186	3813	1.531
40	2581	90		190	4003	1.551
41	2671	90		193	4196	1.571
42	2761	89	90.44	197	4393	1.591
43	2849	89		200	4593	1.612
44	2937	88		203	4796	1.633
45	3024	87		206	5002	1.654
46	3111	86		209	5212	1.675
47	3196	85		212	5423	1.697
48	3280	84		214	5638	1.719
49	3363	83	86.10	217	5855	1.741
50	3445	82		219	6074	1.763
51	3526	81		221	6295	1.785
52	3606	80		223	6518	1.808
53	3684	78		225	6744	1.831
54	3761	77		227	6971	1.854
55	3836	76		229	7200	1.877
56	3911	74	78.20	230	7430	1.900
57	3984	73		232	7661	1.923
58	4055	71		233	7894	1.947
59	4125	70		234	8129	1.970
60	4194	69		235	8364	1.994
61	4261	67		236	8600	2.018
62	4327	66		237	8837	2.042
63	4391	64	68.62	238	9075	2.067
64	4454	63		239	9313	2.091
65	4516	61		239	9553	2.115
66	4576	60		240	9792	2.140
67	4634	59		240	10032	2.165
68	4692	57		240	10273	2.190
69	4747	56		241	10513	2.214
70	4802	55	58.70	241	10754	2.239

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table the values are rounded. This may result in small inaccuracies when using the objectives to calculate other performance statistics.

As-Hatched Performance

Day	Body weight (lb) ¹	Daily gain (lb)	Av. daily gain/week (lb)	Daily intake (lb)	Cum. intake (lb) ²	FCR ³
0	0.096					
1	0.132	0.036			0.024	0.183
2	0.169	0.038		0.033	0.057	0.336
3	0.212	0.042		0.041	0.098	0.463
4	0.260	0.048		0.050	0.148	0.570
5	0.314	0.054		0.059	0.207	0.659
6	0.375	0.061		0.068	0.275	0.735
7	0.442	0.067	0.049	0.078	0.353	0.799
8	0.517	0.075		0.088	0.441	0.854
9	0.598	0.082		0.098	0.538	0.900
10	0.688	0.089		0.108	0.647	0.941
11	0.784	0.097		0.119	0.765	0.976
12	0.889	0.104		0.129	0.895	1.007
13	1.001	0.112		0.140	1.035	1.035
14	1.120	0.119	0.097	0.152	1.187	1.060
15	1.247	0.127		0.163	1.350	1.083
16	1.381	0.134		0.175	1.525	1.104
17	1.523	0.142		0.186	1.711	1.124
18	1.672	0.149		0.198	1.909	1.143
19	1.827	0.155		0.210	2.119	1.161
20	1.989	0.162		0.222	2.342	1.178
21	2.158	0.168	0.148	0.234	2.576	1.195
22	2.332	0.174		0.247	2.823	1.211
23	2.512	0.180		0.259	3.082	1.228
24	2.697	0.185		0.271	3.353	1.244
25	2.888	0.190		0.283	3.637	1.260
26	3.082	0.195		0.296	3.932	1.277
27	3.282	0.199		0.308	4.240	1.293
28	3.485	0.203	0.190	0.320	4.560	1.309
29	3.691	0.207		0.332	4.892	1.326
30	3.901	0.210		0.343	5.235	1.343
31	4.113	0.212		0.355	5.590	1.360
32	4.328	0.215		0.366	5.957	1.377
33	4.545	0.217		0.378	6.334	1.395
34	4.763	0.218		0.389	6.723	1.413
35	4.983	0.220	0.214	0.399	7.122	1.431
36	5.204	0.221		0.410	7.532	1.449
37	5.425	0.221		0.420	7.952	1.467
38	5.647	0.222		0.430	8.382	1.486
39	5.868	0.222		0.440	8.822	1.505
40	6.090	0.221		0.449	9.271	1.524
41	6.310	0.221		0.458	9.729	1.544
42	6.530	0.220	0.221	0.467	10.196	1.563
43	6.749	0.219		0.475	10.671	1.583
44	6.967	0.218		0.483	11.154	1.603
45	7.183	0.216		0.491	11.645	1.623
46	7.397	0.214		0.498	12.143	1.644
47	7.609	0.212		0.505	12.648	1.665
48	7.819	0.210		0.512	13.160	1.686
49	8.027	0.208	0.214	0.518	13.678	1.707
50	8.233	0.205		0.524	14.203	1.728
51	8.436	0.203		0.530	14.733	1.749
52	8.636	0.200		0.536	15.269	1.771
53	8.833	0.197		0.541	15.809	1.793
54	9.028	0.195		0.545	16.355	1.815
55	9.219	0.192		0.550	16.905	1.837
56	9.408	0.189	0.197	0.554	17.459	1.859
57	9.593	0.185		0.558	18.017	1.882
58	9.776	0.182		0.562	18.579	1.904
59	9.955	0.179		0.565	19.144	1.927
60	10.130	0.176		0.568	19.712	1.950
61	10.303	0.172		0.571	20.283	1.973
62	10.472	0.169		0.573	20.856	1.996
63	10.637	0.166	0.176	0.576	21.432	2.019
64	10.800	0.162		0.578	22.010	2.042
65	10.959	0.159		0.580	22.589	2.066
66	11.114	0.156		0.581	23.171	2.089
67	11.267	0.152		0.583	23.753	2.113
68	11.416	0.149		0.584	24.337	2.137
69	11.561	0.146		0.585	24.922	2.161
70	11.704	0.142	0.152	0.586	25.508	2.185

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table the values are rounded. This may result in small inaccuracies when using the objectives to calculate other performance statistics.

Male Performance

Day	Body weight (lb) ¹	Daily gain (lb)	Av. daily gain/week (lb)	Daily intake (lb)	Cum. intake (lb) ²	FCR ³
0	0.096					
1	0.132	0.036			0.022	0.169
2	0.170	0.038		0.031	0.053	0.312
3	0.213	0.043		0.039	0.092	0.433
4	0.262	0.049		0.048	0.141	0.536
5	0.318	0.055		0.058	0.198	0.624
6	0.380	0.062		0.067	0.265	0.699
7	0.449	0.069	0.050	0.077	0.343	0.764
8	0.525	0.076		0.088	0.430	0.819
9	0.609	0.084		0.098	0.529	0.868
10	0.701	0.092		0.109	0.638	0.910
11	0.801	0.100		0.121	0.758	0.947
12	0.909	0.108		0.132	0.890	0.979
13	1.025	0.116		0.144	1.034	1.009
14	1.150	0.124	0.100	0.156	1.190	1.035
15	1.282	0.132		0.168	1.358	1.059
16	1.422	0.140		0.180	1.538	1.082
17	1.570	0.148		0.193	1.731	1.102
18	1.726	0.156		0.206	1.937	1.122
19	1.890	0.164		0.219	2.156	1.140
20	2.061	0.171		0.232	2.387	1.158
21	2.239	0.178	0.156	0.245	2.632	1.176
22	2.423	0.185		0.258	2.890	1.192
23	2.614	0.191		0.271	3.161	1.209
24	2.811	0.197		0.284	3.445	1.225
25	3.014	0.203		0.297	3.742	1.241
26	3.223	0.208		0.311	4.053	1.258
27	3.436	0.213		0.324	4.377	1.274
28	3.654	0.218	0.202	0.337	4.714	1.290
29	3.876	0.222		0.350	5.064	1.306
30	4.102	0.226		0.363	5.426	1.323
31	4.331	0.229		0.375	5.802	1.339
32	4.564	0.232		0.388	6.190	1.356
33	4.799	0.235		0.400	6.590	1.373
34	5.036	0.237		0.412	7.003	1.390
35	5.276	0.239	0.232	0.424	7.427	1.408
36	5.517	0.241		0.436	7.863	1.425
37	5.759	0.242		0.447	8.310	1.443
38	6.001	0.243		0.458	8.768	1.461
39	6.245	0.243		0.469	9.238	1.479
40	6.488	0.243		0.480	9.717	1.498
41	6.731	0.243		0.490	10.207	1.516
42	6.974	0.243	0.243	0.500	10.706	1.535
43	7.216	0.242		0.509	11.215	1.554
44	7.457	0.241		0.518	11.734	1.573
45	7.697	0.240		0.527	12.261	1.593
46	7.936	0.238		0.535	12.796	1.612
47	8.172	0.237		0.544	13.340	1.632
48	8.407	0.235		0.551	13.891	1.652
49	8.639	0.233	0.238	0.559	14.450	1.673
50	8.870	0.230		0.566	15.015	1.693
51	9.097	0.228		0.572	15.588	1.713
52	9.323	0.225		0.579	16.166	1.734
53	9.545	0.222		0.585	16.751	1.755
54	9.764	0.219		0.590	17.342	1.776
55	9.981	0.216		0.596	17.937	1.797
56	10.194	0.213	0.222	0.601	18.538	1.818
57	10.404	0.210		0.605	19.144	1.840
58	10.611	0.207		0.610	19.753	1.862
59	10.815	0.204		0.614	20.367	1.883
60	11.015	0.200		0.618	20.985	1.905
61	11.212	0.197		0.621	21.606	1.927
62	11.405	0.193		0.624	22.230	1.949
63	11.594	0.190	0.200	0.627	22.857	1.971
64	11.780	0.186		0.630	23.487	1.994
65	11.963	0.182		0.632	24.119	2.016
66	12.142	0.179		0.634	24.753	2.039
67	12.317	0.175		0.636	25.390	2.061
68	12.488	0.172		0.638	26.027	2.084
69	12.656	0.168		0.639	26.666	2.107
70	12.821	0.164	0.175	0.640	27.307	2.130

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table the values are rounded. This may result in small inaccuracies when using the objectives to calculate other performance statistics.

Female Performance

Day	Body weight (lb) ¹	Daily gain (lb)	Av. daily gain/week (lb)	Daily intake (lb)	Cum. intake (lb) ²	FCR ³
0	0.096					
1	0.131	0.035			0.026	0.197
2	0.168	0.037		0.035	0.060	0.359
3	0.210	0.042		0.043	0.103	0.493
4	0.257	0.047		0.052	0.155	0.603
5	0.310	0.053		0.060	0.215	0.695
6	0.370	0.059		0.069	0.285	0.771
7	0.435	0.066	0.048	0.078	0.363	0.834
8	0.508	0.073		0.088	0.451	0.888
9	0.588	0.079		0.097	0.548	0.933
10	0.674	0.086		0.107	0.655	0.972
11	0.768	0.094		0.117	0.772	1.006
12	0.868	0.101		0.127	0.899	1.035
13	0.976	0.108		0.137	1.036	1.062
14	1.091	0.115	0.094	0.148	1.184	1.085
15	1.212	0.122		0.158	1.342	1.107
16	1.341	0.128		0.169	1.511	1.127
17	1.476	0.135		0.180	1.691	1.146
18	1.617	0.141		0.191	1.881	1.164
19	1.764	0.147		0.202	2.083	1.181
20	1.918	0.153		0.213	2.296	1.197
21	2.076	0.159	0.141	0.224	2.521	1.214
22	2.240	0.164		0.236	2.756	1.230
23	2.409	0.169		0.247	3.003	1.246
24	2.583	0.174		0.258	3.261	1.263
25	2.761	0.178		0.269	3.531	1.279
26	2.942	0.182		0.281	3.812	1.295
27	3.127	0.185		0.292	4.103	1.312
28	3.316	0.188	0.177	0.303	4.406	1.329
29	3.507	0.191		0.314	4.720	1.346
30	3.700	0.193		0.324	5.044	1.363
31	3.895	0.195		0.335	5.378	1.381
32	4.092	0.197		0.345	5.723	1.399
33	4.291	0.198		0.355	6.079	1.417
34	4.490	0.199		0.365	6.443	1.435
35	4.691	0.200	0.196	0.375	6.818	1.454
36	4.891	0.201		0.384	7.202	1.472
37	5.092	0.201		0.393	7.595	1.492
38	5.292	0.200		0.402	7.996	1.511
39	5.492	0.200		0.410	8.406	1.531
40	5.691	0.199		0.418	8.825	1.551
41	5.889	0.198		0.426	9.251	1.571
42	6.086	0.197	0.199	0.434	9.685	1.591
43	6.282	0.196		0.441	10.126	1.612
44	6.476	0.194		0.448	10.574	1.633
45	6.668	0.192		0.455	11.029	1.654
46	6.858	0.190		0.461	11.490	1.675
47	7.046	0.188		0.467	11.956	1.697
48	7.232	0.186		0.473	12.429	1.719
49	7.415	0.183	0.190	0.478	12.907	1.741
50	7.596	0.181		0.483	13.390	1.763
51	7.774	0.178		0.488	13.878	1.785
52	7.949	0.175		0.492	14.371	1.808
53	8.122	0.173		0.497	14.867	1.831
54	8.291	0.170		0.501	15.368	1.854
55	8.458	0.167		0.504	15.872	1.877
56	8.622	0.164	0.172	0.508	16.380	1.900
57	8.782	0.161		0.511	16.891	1.923
58	8.940	0.158		0.514	17.404	1.947
59	9.094	0.154		0.516	17.920	1.970
60	9.246	0.151		0.519	18.439	1.994
61	9.394	0.148		0.521	18.960	2.018
62	9.539	0.145		0.523	19.482	2.042
63	9.681	0.142	0.151	0.524	20.007	2.067
64	9.819	0.139		0.526	20.533	2.091
65	9.955	0.136		0.527	21.060	2.115
66	10.087	0.132		0.528	21.588	2.140
67	10.217	0.129		0.529	22.117	2.165
68	10.343	0.126		0.530	22.647	2.190
69	10.466	0.123		0.531	23.178	2.214
70	10.587	0.120	0.129	0.531	23.708	2.239

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.


NOTE: In the table the values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.

Carcass Yield

The following diagrams indicate how yields of the major portions change with increasing live weight in each sex. Two types of processing are described: eviscerated yield is broken down into breast meat, thigh and drumstick to represent a portioning operation and into breast meat and leg meat to represent a deboning operation.

Definitions of Terms


Eviscerated %	eviscerated carcass (without neck, abdominal fat and internal organs) as a percentage of live weight.
Breast %	breast meat (without skin and bone) as a percentage of live weight.
Thigh/Drumstick %	whole thigh/drumstick (with skin and bone) as a percentage of live weight.



NOTE: These figures represent dry yield. They do not include any moisture retained during chilling or processing. Carcass component yields will vary among processing plants depending on, for example, type of equipment used and the exact portion(s) being produced.

Definitions of Terms


Eviscerated %	eviscerated carcass (without neck, abdominal fat and internal organs) as a percentage of live weight.
Breast %	breast meat (without skin and bone) as a percentage of live weight.
Thigh/Drumstick %	whole thigh/drumstick (with skin and bone) as a percentage of live weight.



NOTE: These figures represent dry yield. They do not include any moisture retained during chilling or processing. Carcass component yields will vary among processing plants depending on, for example, type of equipment used and the exact portion(s) being produced.

Definitions of Terms


Breast %	breast meat (without skin and bone) as a percentage of live weight.
Leg Meat %	sum of deboned thigh (without skin) and deboned drumstick (without skin) as a percentage of live weight.
Total %	sum of leg meat and breast meat.



NOTE: These figures represent dry yield. They do not include any moisture retained during chilling or processing. Carcass component yields will vary among processing plants depending on, for example, type of equipment used and the exact portion(s) being produced.

Definitions of Terms

Breast %	breast meat (without skin and bone) as a percentage of live weight.
Leg Meat %	sum of deboned thigh (without skin) and deboned drumstick (without skin) as a percentage of live weight.
Total %	sum of leg meat and breast meat.



NOTE: These figures represent dry yield. They do not include any moisture retained during chilling or processing. Carcass component yields will vary among processing plants depending on, for example, type of equipment used and the exact portion(s) being produced.



Aviagen and the Aviagen logo, Ross and the Ross logo, and Yield Plus and the Yield Plus logo are registered trademarks of Aviagen in the US and other countries. All other trademarks or brands are registered by their respective owners.

Privacy Statement: Aviagen collects data to effectively communicate and provide information to you about our products and our business. This data may include your email address, name, business address and telephone number. To view our full Privacy Policy visit [Aviagen.com](https://www.aviagen.com)

© 2021 Aviagen.

Dec 2020